

ESTIMATES OF THE NUMBER OF LINES LYING IN THE LIMIT SET FOR SUBGROUPS OF $PSL(3, \mathbb{C})$

W. BARRERA, A. CANO & J. P. NAVARRETE

ABSTRACT. Given a discrete subgroup $\Gamma \subset PSL(3, \mathbb{C})$, we determine the number of complex lines and complex lines in general position lying in the complement of: maximal regions on which Γ acts properly discontinuously, the Kulkarni's limit set of Γ and the equicontinuity set of Γ . We also provide sufficient conditions to ensure that the equicontinuity region agrees with the Kulkarni's discontinuity region and is the largest set where the group acts properly discontinuously and we provide a description of the respective limit set in terms of the elements of the group.

INTRODUCTION

Given a discrete group $\Gamma \subset PSL(n+1, \mathbb{C})$, $n \neq 3$, is known that Γ may have various maximal sets on which Γ acts properly discontinuously, see [5]. And is not clear how the various extensions of the definition of discontinuity regions, see [5, 6, 8], are related. This article is a continuation of our previous articles, see [1, 2], on which have tried to solve the after told problem in the case $n = 2$. Our argument, in the present article, is based on the existence of lines in the complement of discontinuity regions, see [4], the generalization of the Montel's lemma for Normal families, see [1, 7] and the following estimations, see their respective proofs bellow:

Theorem 0.1. *Let $\Gamma \subset PSL(3, \mathbb{C})$ be a discrete subgroup and $\Omega \subset \mathbb{P}_{\mathbb{C}}^2$ be either $Eq(\Gamma)$, $\Omega(\Gamma)$ or a maximal region where Γ acts properly discontinuously, then $Li(\Omega) = 1, 2, 3$ or ∞ .*

Theorem 0.2. *Let $\Gamma \subset PSL(3, \mathbb{C})$ be a discrete subgroup and $\Omega \subset \mathbb{P}_{\mathbb{C}}^2$ be either $Eq(\Gamma)$, $\Omega(\Gamma)$ or a maximal region where Γ acts properly discontinuously, then $LiG(\Omega) = 1, 2, 3, 4$ or ∞ .*

By means of this results we are able to show:

Theorem 0.3. *Let $\Gamma \subset PSL(3, \mathbb{C})$ be a discrete group. If the number of lines of each maximal set $\Omega \subset \mathbb{P}_{\mathbb{C}}^2$ on which Γ acts properly discontinuously is at least 3, then the Kulkarni's discontinuity region $\Omega(\Gamma)$ agrees with the equicontinuity set of the group Γ , is the largest open set where Γ acts properly*

1991 *Mathematics Subject Classification.* Primary: 32Q45, 37F45; Secondary 22E40.

Key words and phrases. kleinian groups, projective complex plane.

Research partially supported by grants from CNPq.

discontinuously and $\Lambda(\Gamma)$ is the union of complex lines. Moreover, it holds that:

$$\Lambda(\Gamma) = \overline{\bigcup_{\gamma \in \Gamma} \Lambda(\gamma)}.$$

Such theorem extent our previous results in [1].

This article is organized as follows: in section 1 we introduce some terms and notations which will be used along the text. In section 2 we prove Theorems 0.1 and 0.2. In section we prove Theorem 0.3. Finally in section 4, we provide examples which show that our estimates are optimal.

1. PRELIMINARIES AND NOTATIONS

1.1. Projective Geometry. We recall that the complex projective plane $\mathbb{P}_{\mathbb{C}}^2$ is

$$\mathbb{P}_{\mathbb{C}}^2 := (\mathbb{C}^3 \setminus \{0\})/\mathbb{C}^*,$$

where $\mathbb{C}^*$ acts on $\mathbb{C}^3 \setminus \{0\}$ by the usual scalar multiplication. Let $[\] : \mathbb{C}^3 \setminus \{0\} \rightarrow \mathbb{P}_{\mathbb{C}}^2$ be the quotient map. A set $\ell \subset \mathbb{P}_{\mathbb{C}}^2$ is said to be a complex line if $[\ell]^{-1} \cup \{0\}$ is a complex linear subspace of dimension 2. Given $p, q \in \mathbb{P}_{\mathbb{C}}^2$ distinct points, there is a unique complex line passing through p and q , such line will be denoted by $\overleftrightarrow{p, q}$. The set of all the complex lines contained in $\mathbb{P}_{\mathbb{C}}^2$, denoted $Gr(\mathbb{P}_{\mathbb{C}}^2)$, endowed with the topology of the Hausdorff convergence, turns out to be homeomorphic $\mathbb{P}_{\mathbb{C}}^2$.

Consider the action of $\mathbb{Z}_3$ (viewed as the cubic roots of the unity) on $SL(3, \mathbb{C})$ given by the usual scalar multiplication, then

$$PSL(3, \mathbb{C}) = SL(3, \mathbb{C})/\mathbb{Z}_3$$

is a Lie group whose elements are called projective transformations. Let $[[\]] : SL(3, \mathbb{C}) \rightarrow PSL(3, \mathbb{C})$ be the quotient map, $\gamma \in PSL(3, \mathbb{C})$ and $\tilde{\gamma} \in GL(3, \mathbb{C})$, we will say that $\tilde{\gamma}$ is a lift of γ if there is a cubic root τ of $Det(\gamma)$ such that $[[\tau\tilde{\gamma}]] = \gamma$, also, we will use the notation (γ_{ij}) to denote elements in $SL(3, \mathbb{C})$. One can show that $PSL(3, \mathbb{C})$ is a Lie group that acts transitively, effectively and by biholomorphisms on $\mathbb{P}_{\mathbb{C}}^2$ by $[[\gamma]]([w]) = [\gamma(w)]$, where $w \in \mathbb{C}^3 \setminus \{0\}$ and $\gamma \in SL_3(\mathbb{C})$.

Now define the space of the pseudo-projective maps:

$$SP(3, \mathbb{C}) = (M(3, \mathbb{C}) \setminus \{0\})/\mathbb{C}^*,$$

where $\mathbb{C}^*$ acts on $M(3, \mathbb{C}) \setminus \{0\}$ by the usual scalar multiplication. In the sequel $[[\]] : M(3, \mathbb{C}) \setminus \{0\} \rightarrow SP(3, \mathbb{C})$ will denote the quotient map and we will consider $SP(3, \mathbb{C})$ endowed with the quotient topology. Finally, given $\gamma \in SP(3, \mathbb{C})$ we define its kernel by:

$$Ker(\gamma) = [Ker(\tilde{\gamma}) \setminus \{0\}],$$

where $\tilde{\gamma} \in M(3, \mathbb{C})$ is a lift of γ . Clearly $PSL(3, \mathbb{C}) \subset SP(3, \mathbb{C})$ and $\gamma \in PSL(3, \mathbb{C})$ if and only if $Ker(\gamma) = \emptyset$.

1.2. Discontinuous Actions on $\mathbb{P}_{\mathbb{C}}^2$.

Definition 1.1. Let $\Gamma \subset PSL(3, \mathbb{C})$ be a discrete group. An open non-empty set $\Omega \subset \mathbb{P}_{\mathbb{C}}^2$ is said to be a region where Γ acts properly discontinuously if $\Gamma\Omega = \Omega$ and for each compact set $K \subset \Omega$ the set $\{\gamma \in \Gamma \mid \gamma(K) \cap K\}$ is finite. Set

$$Dis(\Gamma) = \{\Omega \subset \mathbb{P}_{\mathbb{C}}^2 : \Omega \neq \emptyset \text{ \& } \Gamma \text{ acts properly discontinuously on } \Omega\}.$$

The group Γ is said to be Kleinian if $Dis(\Gamma) \neq \emptyset$.

1.2.1. *The equicontinuity set.* Recall that the *equicontinuity set* for a family $\mathcal{F}$ of endomorphisms of $\mathbb{P}_{\mathbb{C}}^2$, denoted $Eq(\mathcal{F})$, is defined to be the set of points $z \in \mathbb{P}_{\mathbb{C}}^2$ for which there is an open neighborhood U of z such that $\{f|_U : f \in \mathcal{F}\}$ is a normal family.

Definition 1.2. Let $\Gamma \subset PSL(3, \mathbb{C})$ be a discrete group, then we define:

$$Lim(\Gamma) = \{g \in SP(3, \mathbb{C}) : g \text{ is a cluster point of } \Gamma\}.$$

Theorem 1.3 (See [1]). *If $\Gamma \subset PSL(3, \mathbb{C})$ is a discrete group, then:*

- (i) *The equicontinuity region of Γ is a region where Γ acts properly discontinuously;*
- (ii) *The equicontinuity region is described as follows:*

$$Eq(\Gamma) = \mathbb{P}_{\mathbb{C}}^2 \setminus \left(\bigcup_{\gamma \in Lim(\Gamma)} Ker(\gamma) \right);$$

- (iii) *If U is a Γ -invariant open subset of $\mathbb{P}_{\mathbb{C}}^2$ such that $\mathbb{P}_{\mathbb{C}}^2 \setminus U$ contains at least three complex lines in general position, then $U \subset Eq(\Gamma)$.*

1.2.2. *The Discontinuity Region in the Kulkarni's sense.* Given $\Gamma \subset PSL(3, \mathbb{C})$ a discrete group we define following Kulkarni, see [8]: the set $L_0(\Gamma)$ as the closure of the points in $\mathbb{P}_{\mathbb{C}}^2$ with infinite isotropy group. The set $L_1(\Gamma)$ as the closure of the set of cluster points of Γz where z runs over $\mathbb{P}_{\mathbb{C}}^2 \setminus L_0(\Gamma)$. The set $L_2(\Gamma)$ as the closure of cluster points of ΓK where K runs over all the compact sets in $\mathbb{P}_{\mathbb{C}}^2 \setminus (L_0(\Gamma) \cup L_1(\Gamma))$. The *Limit Set in the sense of Kulkarni* for Γ is defined as:

$$\Lambda(\Gamma) = L_0(\Gamma) \cup L_1(\Gamma) \cup L_2(\Gamma).$$

The *Discontinuity Region in the sense of Kulkarni* of Γ is defined as:

$$\Omega(\Gamma) = \mathbb{P}_{\mathbb{C}}^2 \setminus \Lambda(\Gamma).$$

Clearly the discontinuity region $\Omega(\Gamma)$ contains $Eq(\Gamma)$, see [1], and is a region where Γ acts properly discontinuously, see [8].

2. COUNTING LINES

Definition 2.1. Let $\Omega \subset \mathbb{P}_{\mathbb{C}}^2$ be a non-empty open set. Let us define:

- (i) The lines in general position outside Ω as:

$$LG(\Omega) = \left\{ \mathcal{L} \subset Gr_1(\mathbb{P}_{\mathbb{C}}^2) \mid \text{The lines in } \mathcal{L} \text{ are in general position \& } \bigcup \mathcal{L} \subset \mathbb{P}_{\mathbb{C}}^2 \setminus \Omega \right\};$$

- (ii) The number of lines in general position outside Ω as:

$$LiG(\Omega) = \max(\{card(\mathcal{L}) : \mathcal{L} \in LG(\Omega)\}),$$

where $card(C)$ denotes the number of elements contained in C .

- (iii) Given $\mathcal{L} \in LG(\Omega)$ and $v \in \bigcup \mathcal{L}$, we will say that v is a vertex for $\mathcal{L}$ if there are $\ell_1, \ell_2 \in \mathcal{L}$ distinct lines and an infinite set $\mathcal{C} \subset Gr_1(\mathbb{P}_{\mathbb{C}}^2)$ such that $\ell_1 \cap \ell_2 \cap (\bigcap \mathcal{C}) = \{v\}$ and $\bigcup \mathcal{C} \subset \mathbb{P}_{\mathbb{C}}^2 \setminus \Omega$.

Corollary 2.2. Let $\Gamma \subset PSL(3, \mathbb{C})$ be a discrete group and $\Omega \subset \mathbb{P}_{\mathbb{C}}^2$ be a maximal discontinuity region for the action of Γ if $LiG(\Omega(\Gamma)) \leq 2$, then $LiG(\Omega) \leq 2$.

Proof. On the contrary, let us assume that there is a maximal region $\Omega \subset \mathbb{P}_{\mathbb{C}}^2$ where Γ acts properly discontinuously and $LiG(\Omega) \geq 3$, thus $\Omega \subset Eq(\Gamma) \subset \Omega(\Gamma)$. Thus $\Omega = \Omega(\Gamma)$, which is a contradiction. $\square$

Definition 2.3. Let $\Gamma \subset PSL(3, \mathbb{C})$ be a group and $p \in \mathbb{P}_{\mathbb{C}}^2$ such that $\Gamma p = p$, then define $\Pi = \Pi_{p, \ell, \Gamma} : \Gamma \longrightarrow Bihol(\ell)$ given by $\Pi(g)(x) = \pi(g(x))$ where $\pi = \pi_{p, \ell} : \mathbb{P}_{\mathbb{C}}^2 - \{p\} \longrightarrow \ell$ is given by

$$\pi(x) = \overleftrightarrow{xp} \cap \ell.$$

Also, if $p = e_1$ define $\lambda_{\infty, \Gamma} : \Gamma \rightarrow \mathbb{C}$ by

$$\lambda_{\infty, \Gamma}([\gamma]) = \gamma_{11}^3.$$

Clearly, $\Pi_{p, \ell, \Gamma}$ and $\lambda_{\infty, \Gamma}$ are well defined group morphisms.

Lemma 2.4. Let $\Gamma \subset PSL(3, \mathbb{C})$ be a discrete group and $\Omega \subset \mathbb{P}_{\mathbb{C}}^2$ be an open Γ -invariant set such that $LiG(\Omega) = 2$, then there is a point $p \in \mathbb{P}_{\mathbb{C}}^2$ such that $\Gamma p = p$.

Proof. Let $\mathcal{L} = \{\ell \in Gr(\mathbb{P}_{\mathbb{C}}^2) : \ell \in \mathbb{P}_{\mathbb{C}}^2 \setminus \Omega\}$. Since $LiG(\Omega) = 2$, it follows that $\bigcap \mathcal{L}$ contains exactly one point, namely p_0 . Clearly $\Gamma \mathcal{L} = \mathcal{L}$ and in consequence $\Gamma p = p$. $\square$

Lemma 2.5. Let $\Gamma \subset PSL(3, \mathbb{C})$ be a discrete group and $\Omega \subset \mathbb{P}_{\mathbb{C}}^2$ be an open Γ -invariant set such that $L_0(\Gamma) \subset \mathbb{P}_{\mathbb{C}}^2 \setminus \Omega$. If $LiG(\Omega) = 2$ and $3 \leq Li(\Omega) < \infty$, then:

- (i) If $\ell \in Gr(\mathbb{P}_{\mathbb{C}}^2)$ does not contain p , then $\Pi_{p, \ell}(\Gamma)$ is finite.
- (ii) It holds that $Ker(\Pi)$ is a normal subgroup of finite index.

- (iii) *By conjugating with a projective transformation, if it is necessary, it follows that each $\gamma \in Ker(\Pi)$ with infinite order has a lift $\tilde{\gamma} \in SL(3, \mathbb{C})$ given by:*

$$\tilde{\gamma} = \begin{pmatrix} 1 & b & c \\ 0 & 1 & 0 \\ 0 & 0 & 1 \end{pmatrix}.$$

- (iv) *By conjugating with a projective transformation, if it is necessary, it follows that each $\gamma \in Ker(\Pi)$ with infinite order has a lift $\tilde{\gamma} \in SL(3, \mathbb{C})$ given by:*

$$\tilde{\gamma} = \begin{pmatrix} 1 & b & 0 \\ 0 & 1 & 0 \\ 0 & 0 & 1 \end{pmatrix}.$$

- (v) *By conjugating with a projective transformation, if it is necessary, it follows that the group $\lambda_{\infty, \Gamma}(Ker(\Pi))$ is finite.*

- (vi) *There is $\ell \in Gr(\mathbb{P}_{\mathbb{C}}^2)$ such that $Eq(\Gamma) = \mathbb{P}_{\mathbb{C}}^2 \setminus \ell$.*

Proof. Set $\mathcal{L} = \{\ell \in Gr(\mathbb{P}_{\mathbb{C}}^2) : \ell \in \mathbb{P}_{\mathbb{C}}^2 \setminus \Omega\}$ and $n_0 = Card(\mathcal{L})$.

Let us prove (i). Since $\Gamma \cup \mathcal{L} = \bigcup \mathcal{L}$ it follows that

$$\Pi(\Gamma)\pi\left(\bigcup \mathcal{L} \setminus \{p_0\}\right) = \pi\left(\bigcup \mathcal{L} \setminus \{p_0\}\right).$$

In consequence $\pi\left(\bigcup \mathcal{L} \setminus \{p_0\}\right)$ is a $\Pi(\Gamma)$ -invariant set whose cardinality is n_0 . Thus

$$\Gamma_0 = \bigcap_{x \in \pi\left(\bigcup \mathcal{L} \setminus \{p_0\}\right)} Isot(x, \Pi(\Gamma)),$$

is a normal subgroup of $\Pi(\Gamma)$ with finite index. Moreover, every element in $\pi\left(\bigcup \mathcal{L} \setminus \{p_0\}\right)$ is fixed by Γ_0 . Since $\pi\left(\bigcup \mathcal{L} \setminus \{p_0\}\right)$ contains more than 3 elements we conclude that Γ_0 consist exactly on the identity. Therefore $\Pi(\Gamma)$ is finite.

Let us prove (ii). From the previous claim it follows that $Ker(\Pi)$ is a normal subgroup of Γ with finite index.

Let us prove (iii). Let $\gamma \in Ker(\Pi)$ and $\tilde{\gamma} \in SL(3, \mathbb{C})$ a lift of γ , thus

$$\tilde{\gamma} = \begin{pmatrix} a^{-2} & b & c \\ 0 & a & 0 \\ 0 & 0 & a \end{pmatrix}$$

where $a \in \mathbb{C}^*$ and $b, c \in \mathbb{C}$. if $\tilde{\gamma}$ is diagonalizable, there are $v_1, v_2 \in \mathbb{C}^3$ such that $\{e_1, v_1, v_2\}$ is an eigenbasis of $\tilde{\gamma}$ whose respective eigenvalues are $\{a^{-2}, a, a\}$. In consequence $\ell = \overleftrightarrow{v_1, v_2} \subset Fix(\gamma) \subset \mathbb{P}_{\mathbb{C}}^2 \subset \Omega$ and $e_1 \notin \ell$, which is a contradiction. In consequence $\tilde{\gamma}$ is non diagonalizable and the claim

follows from the Jordan's normal form theorem.

By the previous claim we can assume that there is an element $\gamma_0 \in \text{Ker}(\Pi)$, such that γ_0 has a lift $\tilde{\gamma}_0 \in SL(3, \mathbb{C})$ which is given by:

$$\tilde{\gamma}_0 = \begin{pmatrix} 1 & 1 & 0 \\ 0 & 1 & 0 \\ 0 & 0 & 1 \end{pmatrix}$$

Let us prove (iv). If this is not the case, there is $\gamma_1 \in \text{Ker}(\Pi)$ which has a lift $\tilde{\gamma}_1 \in SL(3, \mathbb{C})$ given by:

$$\tilde{\gamma}_1 = \begin{pmatrix} 1 & a & b \\ 0 & 1 & 0 \\ 0 & 0 & 1 \end{pmatrix}.$$

An inductive argument shows:

$$\tilde{\gamma}_0^m \tilde{\gamma}_1 = \begin{pmatrix} 1 & a+n & b \\ 0 & 1 & 0 \\ 0 & 0 & 1 \end{pmatrix}.$$

A straightforward calculation shows that

$$\ell_m = \overleftarrow{[e_1], [0 : -b : b+n]} \subset \text{Fix}(\gamma_0^m \gamma_1)$$

and $\ell_m \neq \ell_n$, whenever $n \neq m$. That is $\ell_m \in \mathcal{L}$, which is a contradiction. Which concludes the proof of the claim.

Let us prove (v). On the contrary, let us assume that $\lambda_{\infty, \Gamma}(\text{Ker}(\Pi))$ is infinite, thus there is a sequence $\lambda_m^3 \subset \lambda_{\infty, \Gamma}(\text{Ker}(\Pi))$ of distinct elements such that $\lambda_m^{1/2} \rightarrow 1$. For each $m \in \mathbb{N}$ let $\gamma_m \in \text{Ker}(\Pi)$ be such that $\lambda_{\infty, \Gamma}(\gamma_m) = \lambda_m^3$. Therefore γ_m has a lift $\tilde{\gamma}_m \in SL(3, \mathbb{C})$ given by:

$$\tilde{\gamma} = \begin{pmatrix} \lambda_m & a_m & b_m \\ 0 & \lambda_m^{-1/2} & 0 \\ 0 & 0 & \lambda_m^{-1/2} \end{pmatrix}.$$

where $a_m, b_m \in \mathbb{C}$. Now a simple calculation shows:

$$\tilde{\gamma}_m^{-1} \tilde{\gamma}_0 \tilde{\gamma}_m = \begin{pmatrix} 1 & \lambda_m^{-3/2} & 0 \\ 0 & 1 & 0 \\ 0 & 0 & 1 \end{pmatrix} \xrightarrow{m \rightarrow \infty} \begin{pmatrix} 1 & 1 & 0 \\ 0 & 1 & 0 \\ 0 & 0 & 1 \end{pmatrix}.$$

Which is a contradiction, since Γ is discrete.

Let us prove (vi). By the previous claims, it follows that each element $\gamma \in \text{Ker}(\lambda_{\infty, \text{Ker}(\Pi)})$ has a lift $\tilde{\gamma} \in SL(3, \mathbb{C})$ which is given by:

$$\tilde{\gamma} = \begin{pmatrix} 1 & a & 0 \\ 0 & 1 & 0 \\ 0 & 0 & 1 \end{pmatrix}.$$

It follows that $Eq(Ker(\lambda_{\infty, Ker(\Pi)})) = \mathbb{P}_{\mathbb{C}}^2 \setminus \overleftrightarrow{e_1, e_3}$. Which concludes the proof of the claim. $\square$

Definition 2.6. Let $\Gamma \subset PSL(3, \mathbb{C})$ be a discrete group, let us define $C(\Gamma)$ as the set:

$$C(\Gamma) = \overline{\bigcup_{\gamma \in \Gamma} \Lambda(\gamma)}.$$

Clearly $C(\Gamma) \subset \mathbb{P}_{\mathbb{C}}^2 \setminus Eq(\Gamma)$ is a non empty, Γ -invariant closed set, see [9], [4].

Proposition 2.7. *Let $\Gamma \subset PSL(3, \mathbb{C})$ be a discrete subgroup and $\Omega \subset \mathbb{P}_{\mathbb{C}}^2$ be either $Eq(\Gamma)$, $\Omega(\Gamma)$, $\mathbb{P}_{\mathbb{C}}^2 \setminus C(\Gamma)$ or a maximal region where Γ acts properly discontinuously. If $LiG(\Omega) = 2$ and $Li(\Omega) < \infty$, then $LiG(\Omega) = 2$.*

Proof. If Ω is either $Eq(\Gamma)$, $\Omega(\Gamma)$ or $\mathbb{P}_{\mathbb{C}}^2 \setminus C(\Gamma)$ and the assertion does not follows, then previous lemma yields that $\mathbb{P}_{\mathbb{C}}^2 \setminus (\Omega)$ is a complex line, which is a contradiction.

Finally, if Ω is a maximal region where Γ acts properly discontinuously and $Li(\Omega) > 2$, then parts (vi) and (??) of previous lemma yields $L_0(\Gamma) = \mathbb{P}_{\mathbb{C}}^2 \setminus Eq(\Gamma)$ is a complex line. Thus $\Omega \subset Eq(\Gamma)$, which is a contradiction. $\square$

Proposition 2.8. *Let $\Gamma \subset PSL(3, \mathbb{C})$ be a discrete group and $\Omega \subset \mathbb{P}_{\mathbb{C}}^2$ be an open Γ -invariant set such that $L_0(\Gamma) \subset \mathbb{P}_{\mathbb{C}}^2 \setminus \Omega$. If $LiG(\Omega) = 3$ and $3 < Li(\Omega) < \infty$, then there is $\ell \in Gr(\mathbb{P}_{\mathbb{C}}^2)$ and $p \in \mathbb{P}_{\mathbb{C}}^2 \setminus \ell$ such that $Eq(\Gamma) = \mathbb{P}_{\mathbb{C}}^2 \setminus (\ell \cup \{p\}) = \mathbb{P}_{\mathbb{C}}^2 \setminus L_0(\Gamma)$.*

Proof. On the contrary, let us assume that $n_0 = Li(\Omega) \geq 3$. Let $\mathcal{L} = \{\ell \in Gr(\mathbb{P}_{\mathbb{C}}^2) | \ell \in \mathbb{P}_{\mathbb{C}}^2 \setminus \Omega\}$. Let $\Gamma_0 = \bigcap_{\ell \in \mathcal{L}} Isot(\ell, \cdot)$. Since $\mathcal{L}$ is a finite set, we conclude that Γ_0 is a normal subgroup of Γ with finite index. Since $LiG(\Omega) = 3$, and after conjugating by a projective transformation if it is necessary, we can assume that $X = \overleftrightarrow{[e_1], [e_2]}$, $Y = \overleftrightarrow{[e_3], [e_2]}$, $Z = \overleftrightarrow{[e_1], [e_3]}$ $\in \mathcal{L}$. Thus every element $\gamma \in \tau\Gamma_0\tau^{-1}$ has a lift in $\tilde{\gamma} \in SL(3, \mathbb{C})$ given by:

$$\tilde{\gamma} = \begin{pmatrix} \gamma_{11} & 0 & 0 \\ 0 & \gamma_{22} & 0 \\ 0 & 0 & \gamma_{33} \end{pmatrix}.$$

where $\gamma_{11}\gamma_{22}\gamma_{33} = 1$. Let $\ell \in \mathcal{L} \setminus \{X, Y, Z\}$ be a complex line, then either $[e_1] \in \ell$, $[e_2] \in \ell$ or $[e_3] \in \ell$. Without loss of generality let us assume that $e_1 \in \ell$. Set $\Pi = \Pi_{[e_1], Y}$ and $\pi = \pi_{[e_1], Y}$. Thus $[e_2], [e_3], \pi(\ell \setminus [e_1]) \in \bigcap_{\gamma \in \Pi(\Gamma_0)} Fix(\gamma)$ and $[e_2] \neq \pi(\ell \setminus [e_1]) \neq [e_3]$. It falls out that $\Pi(\Gamma_0) = \{Id\}$. Therefore, for each $\gamma \in \Gamma_0$ there is $\gamma_{11} \in \mathbb{C}^*$ such that $\tilde{\gamma} \in SL(3, \mathbb{C})$ given by:

$$(2.1) \quad \tilde{\gamma} = \begin{pmatrix} \gamma_{11}^2 & 0 & 0 \\ 0 & \gamma_{11}^{-1} & 0 \\ 0 & 0 & \gamma_{11}^{-1} \end{pmatrix},$$

is a lift of γ . Therefore

$$(2.2) \quad \begin{aligned} Y \cup \{[e_1]\} &\subset \bigcap_{\gamma \in \Pi(\Gamma_0)} \text{Fix}(\gamma) \\ Eq(\Gamma_0) &= \mathbb{P}_{\mathbb{C}}^2 \setminus (Y \cup \{[e_1]\}). \end{aligned}$$

□

Proposition 2.9. *Let $\Gamma \subset PSL(3, \mathbb{C})$ be a discrete subgroup and $\Omega \subset \mathbb{P}_{\mathbb{C}}^2$ be either $Eq(\Gamma)$, $\Omega(\Gamma)$, $\mathbb{P}_{\mathbb{C}}^2 \setminus C(\Gamma)$ or a maximal region where Γ acts properly discontinuously. If $LiG(\Omega) = 3$ and $Li(\Omega) < \infty$, then $LiG(\Omega) = 3$.*

Proof. If the assertion does not follow, then previous lemma yields that $Li(\Omega) = 1$, which is a contradiction. □

Proposition 2.10. *Let $\Gamma \subset PSL(3, \mathbb{C})$ be a discrete subgroup and $\Omega \subset \mathbb{P}_{\mathbb{C}}^2$ be either $Eq(\Gamma)$, $\Omega(\Gamma)$, $\mathbb{P}_{\mathbb{C}}^2 \setminus C(\Gamma)$ or a maximal region where Γ acts properly discontinuously. If $LiG(\Omega) \geq 4$, then $Li(\Omega)$ is infinite.*

Proof. On the contrary, let us assume that $Li(\Omega) < \infty$. Set $\mathcal{L} = \{\ell \in Gr(\mathbb{P}_{\mathbb{C}}^2) : \ell \subset \mathbb{P}_{\mathbb{C}}^2 \setminus \Omega\}$ and $\Gamma_0 = \bigcap_{\ell \in \mathcal{L}} Isot(\ell, \Gamma)$. Since $LiG(\Omega) \geq 4$, it falls out that each $\gamma \in \Gamma_0$ fixes at least four lines in general position. Thus $\Gamma_0 = \{Id\}$. Since Γ is a subgroup of Γ with finite index, we conclude that Γ is finite. Which is a contradiction. □

Is worth nothing to check that the following theorem follows from the following Lemma.

Lemma 2.11 (See [4] & [10]). *Let $\Gamma \subset PSL(3, \mathbb{C})$ be a discrete subgroup and $\Omega \subset \mathbb{P}_{\mathbb{C}}^2$ be either $Eq(\Gamma)$, $\Omega(\Gamma)$, $\mathbb{P}_{\mathbb{C}}^2 \setminus C(\Gamma)$ or a maximal region where Γ acts properly discontinuously, then Ω contains a complex line.*

Theorem 2.12. *Let $\Gamma \subset PSL(3, \mathbb{C})$ be a discrete subgroup and $\Omega \subset \mathbb{P}_{\mathbb{C}}^2$ be either $Eq(\Gamma)$, $\Omega(\Gamma)$, $\mathbb{P}_{\mathbb{C}}^2 \setminus C(\Gamma)$ or a maximal region where Γ acts properly discontinuously, then $Li(\Omega) = 1, 2, 3$ or ∞ .*

Lemma 2.13. *Let $\Gamma \subset PSL(3, \mathbb{C})$ be a discrete group and $\ell \in Gr(\mathbb{P}_{\mathbb{C}}^2)$ be Γ -invariant such that the action of Γ restricted to ℓ is trivial, then:*

- (i) *If there is an element in Γ with infinite order with a diagonalizable lift, then there is a discrete subgroup $G \subset \mathbb{C}^*$ such that Γ is conjugated to:*

$$\Gamma_G = \left\{ \begin{pmatrix} a & 0 & 0 \\ 0 & a & 0 \\ 0 & 0 & a^{-2} \end{pmatrix} : a \in G \right\}.$$

- (ii) *If Γ does not contain an element with infinite order and with a diagonalizable lift, then, after conjugating with a projective transformation if it is necessary, each element $\gamma \in \Gamma$ has a lift $\tilde{\gamma} \in SL(3, \mathbb{C})$ given by:*

$$\tilde{\gamma} = \begin{pmatrix} a & 0 & b \\ 0 & a & c \\ 0 & 0 & a^{-2} \end{pmatrix},$$

where $|a| = 1$.

Proof. Let us prove (i). After conjugating with a projective transformation, if it is necessary, we may assume that $\ell = \overleftrightarrow{e_1, e_2}$ and there is $\gamma_0 \in \Gamma$ with a lift $\tilde{\gamma}_0 \in SL(3, \mathbb{C})$ given by:

$$\tilde{\gamma}_0 = \begin{pmatrix} a & 0 & 0 \\ 0 & a & 0 \\ 0 & 0 & a^{-2} \end{pmatrix},$$

where $|a| < 1$. On the other hand, since each element $\gamma \in \Gamma$ has a lift $\tilde{\gamma} \in SL(3, \mathbb{C})$ given by:

$$\tilde{\gamma} = \begin{pmatrix} \gamma_{11} & 0 & \gamma_{13} \\ 0 & \gamma_{11} & \gamma_{23} \\ 0 & 0 & \gamma_{11}^{-2} \end{pmatrix},$$

Set $\Pi_1 = \Pi_{e_1, \overleftrightarrow{e_2, e_3}}$, $\pi_1 = \Pi_{e_1, \overleftrightarrow{e_2, e_3}}$ and $\Pi_2 = \Pi_{e_2, \overleftrightarrow{e_1, e_3}}$, $\pi_2 = \Pi_{e_2, \overleftrightarrow{e_1, e_3}}$. Clearly $\text{Ker}(\Pi_j)$, $j \in \{1, 2\}$, is trivial.

Is straightforward to check that the following claim proves the assertion.

Claim 1.- For each $\tau \in \Gamma \setminus \{Id\}$ it follows that $\text{Fix}(\Pi_j(\tau)) = \text{Fix}(\Pi_j(\gamma_0))$. If this is not the case, there are $j_0 \in \{1, 2\}$ and $\tau_0 \in \Gamma$ such that $\text{Fix}(\Pi_{j_0}(\tau_0)) \cap \text{Fix}(\Pi_{j_0}(\gamma_0))$ is a single point. In consequence $\Pi_{j_0}(\tau_0 \gamma_0 \tau_0^{-1} \gamma_0^{-1})$ is parabolic. Without loss of generality, let us assume that $j_0 = 1$, thus $\kappa = \tau_0 \gamma_0 \tau_0^{-1} \gamma_0^{-1}$ has a lift $\tilde{\kappa} \in SL(3, \mathbb{C})$ given by:

$$\tilde{\kappa} = \begin{pmatrix} 1 & 0 & \kappa_{13} \\ 0 & 1 & \kappa_{23} \\ 0 & 0 & 1 \end{pmatrix},$$

where $\kappa_{13} \neq 0$. Finally, a simple calculation shows:

$$\tilde{\gamma}_0^m \tilde{\kappa} \tilde{\gamma}_0^{-m} = \begin{pmatrix} 1 & 0 & a^{-m} \kappa_{13} \\ 0 & 1 & a^{-m} \kappa_{23} \\ 0 & 0 & 1 \end{pmatrix} \xrightarrow{m \rightarrow \infty} \begin{pmatrix} 1 & 0 & 0 \\ 0 & 1 & 0 \\ 0 & 0 & 1 \end{pmatrix},$$

which is a contradiction since Γ is discrete.

Let us prove (ii). After conjugating with a projective transformation, if it is necessary, we may assume that $\ell = \overleftrightarrow{e_1, e_2}$. In consequence each element $\gamma \in \Gamma$ has a lift given by:

$$\tilde{\gamma} = \begin{pmatrix} \gamma_{11} & 0 & \gamma_{13} \\ 0 & \gamma_{11} & \gamma_{23} \\ 0 & 0 & \gamma_{11}^{-2} \end{pmatrix}.$$

Set $\Pi_1 = \Pi_{e_1, \overleftrightarrow{e_2, e_3}}$, $\pi_1 = \Pi_{e_1, \overleftrightarrow{e_2, e_3}}$, $\Pi_2 = \Pi_{e_2, \overleftrightarrow{e_1, e_3}}$, $\pi_2 = \Pi_{e_2, \overleftrightarrow{e_1, e_3}}$ and $\lambda_\infty : (\Gamma) \rightarrow \mathbb{C}^*$ given by $\lambda_\infty(\gamma) = \gamma_{11}^3$.

Clearly, the assertion yields from the following claim:

Claim 2.- $\lambda_\infty(\Gamma) \subset \mathbb{S}^1$. On the contrary, let us assume that there is $\gamma_0 \in \Gamma$ such that $|\lambda_\infty(\gamma_0)| \neq 1$. In such case:

$$\text{Fix}(\Pi_1(\gamma_0)) = \{p_1 = [0 : \gamma_{23}^{(0)} : 1 - \gamma_{11}^{(0)}], [e_2]\}$$

$$\text{Fix}(\Pi_2(\gamma_0)) = \{p_2 = [\gamma_{13}^{(0)} : 0 : 1 - \gamma_{11}^{(0)}], [e_1]\}.$$

Set $\ell_1 = \overrightarrow{p_1, e_1}$ and $\ell_2 = \overrightarrow{p_2, e_2}$. Thus $\gamma_0(\ell_1) = \ell_1$ and $\gamma_0(\ell_2) = \ell_2$. That is the unique point $*$ in $\ell_1 \cap \ell_2$ is fixed by γ_0 and $*$ $\in \mathbb{P}_{\mathbb{C}}^2 \setminus (\ell_1 \cup \ell_2)$. Therefore γ_0 is diagonalizable. Finally, since λ_∞ is a group morphism, we conclude that γ_0 has infinite order. Which contradicts the initial assumption. $\square$

Now it follows easily that:

Corollary 2.14. *Let $\Gamma \subset PSL(3, \mathbb{C})$ be a discrete group and $\ell \in Gr(\mathbb{P}_{\mathbb{C}}^2)$ be an invariant line under Γ such that the action of Γ restricted to ℓ is trivial, then there is a point $p \in \mathbb{P}_{\mathbb{C}}^2$ such that $\text{Eq}(\Gamma) = \mathbb{P}^2 \setminus (\ell \cup \{p\})$.*

Lemma 2.15. *Let $\Gamma \subset PSL(3, \mathbb{C})$ be a discrete group such that each element $\gamma \in \Gamma$ has a lift $\tilde{\gamma} \in GL(3, \mathbb{C})$ given by:*

$$\tilde{\gamma} = \begin{pmatrix} \gamma_{11} & 0 & 0 \\ 0 & \gamma_{22} & \gamma_{23} \\ 0 & \gamma_{32} & \gamma_{33} \end{pmatrix},$$

then $\text{LiG}(\text{Eq}(\Gamma)) \leq 3$.

Proof. Let $(\gamma_m) \subset \Gamma$ be a sequence of distinct, thus for each $m \in \mathbb{N}$ there is $\tilde{\gamma}_m \in GL(3, \mathbb{C})$ a lift of γ_m which is given by:

$$\tilde{\gamma}_m = \begin{pmatrix} \gamma_{11}^{(m)} & 0 & 0 \\ 0 & \gamma_{22}^{(m)} & \gamma_{23}^{(m)} \\ 0 & \gamma_{32}^{(m)} & \gamma_{33}^{(m)} \end{pmatrix}$$

where $\max\{|\gamma_{11}^{(m)}|, |\gamma_{22}^{(m)}|, |\gamma_{33}^{(m)}|, |\gamma_{23}^{(m)}|, |\gamma_{32}^{(m)}|\} = 1$. Since the sequences $(\gamma_{11}^{(m)}), (\gamma_{22}^{(m)}), (\gamma_{33}^{(m)}), (\gamma_{23}^{(m)}), (\gamma_{32}^{(m)})$ are bounded, there is a sequence $(k_m) \subset (m)$ and $\gamma_{11}, \gamma_{22}, \gamma_{33}, \gamma_{23}, \gamma_{32} \in \mathbb{C}$ such that

$$\tilde{\gamma}_{k_m} \xrightarrow{m \rightarrow \infty} \begin{pmatrix} \gamma_{11} & 0 & 0 \\ 0 & \gamma_{22} & \gamma_{23} \\ 0 & \gamma_{32} & \gamma_{33} \end{pmatrix} = \tilde{\gamma}$$

where $\gamma_{11}(\gamma_{22}\gamma_{33} - \gamma_{32}\gamma_{23}) = 0$. In consequence $\text{Ker}([\tilde{\gamma}])$ is either $[e_1]$, a line passing trough $[e_1], [e_2], [e_3]$ or a point in $[e_2], [e_3]$. Now, the conclusion follows easily. $\square$

Proposition 2.16. *Let $\Gamma \subset PSL(3, \mathbb{C})$ be a discrete group and $\Omega \subset \mathbb{P}_{\mathbb{C}}^2$ be a discontinuity region for the action of Γ which is either $\text{Eq}(\Gamma)$, $\Omega(\Gamma)$ or a maximal discontinuity region. If $4 \leq \text{LiG}(\Omega) < \infty$, then for each $\mathcal{L} \in GL(\Omega)$ with $\text{Card}(\mathcal{L} = \text{LiG}(\Omega))$ it falls out that:*

- (i) *There is at least one vertex of $\mathcal{L}$;*
- (ii) *If $v \in \mathbb{P}_{\mathbb{C}}^2$ is a vertex of $\mathcal{L}$, then $Isot(v, \Gamma)$ is a subgroup of Γ with finite index;*
- (iii) *There are at least two vertexes of $\mathcal{L}$;*
- (iv) *The vertexes of $\mathcal{L}$ lie in a complex line;*
- (v) *There are at most two vertexes of $\mathcal{L}$;*
- (vi) *If $\ell \in Gr(\mathbb{P}_{\mathbb{C}}^2)$ does not contains a vertex of $\mathcal{L}$, then its orbit under Γ is infinite.*

Proof. Let us prove part (i). Since $LiG(\Omega) \geq 4$ we conclude that $Lin(\Omega) = \infty$. On the other hand, since $LinG(\Omega) < \infty$, we deduce that there is $p_0 \in \mathbb{P}_{\mathbb{C}}^2$, $\ell_1, \ell_2 \in \mathcal{L}$ and a infinite set $\mathcal{C}_0$ such that $\{p_0\} = \bigcap \mathcal{C}_0 = \ell_1 \cap \ell_2$ and $\bigcup \mathcal{C}_0 \subset \mathbb{P}_{\mathbb{C}}^2 \setminus \Omega$.

Let us prove (ii). Let $p_0 \in \tilde{\mathcal{P}}(\mathcal{L})$, then there are $\ell_1, \ell_2 \in \mathcal{L}$ and a infinite set $\mathcal{C}_0$ such that $\{p_0\} = \bigcap \mathcal{C}_0 = \ell_1 \cap \ell_2$ and $\bigcup \mathcal{C}_0 \subset \mathbb{P}_{\mathbb{C}}^2 \setminus \Omega$. Let $\gamma \in \Gamma$, then there is $p_1 \in \mathbb{P}_{\mathbb{C}}^2$, $\tilde{\ell}_1, \tilde{\ell}_2 \in \mathcal{L}$ and a infinite set $\mathcal{C}_1 \subset \mathcal{C}_0$ such that $\{p_1\} = \bigcap \gamma(\mathcal{C}_0) = \tilde{\ell}_1 \cap \tilde{\ell}_2$. In consequence $\gamma(p_0) = p_1$. Which proves the assertion.

Let us prove (iii). Let $p_0 \in \mathbb{P}_{\mathbb{C}}^2$ be a vertex of $\mathcal{L}$, thus there are $\ell_3, \ell_4 \in \mathcal{L}$ such that $p_0 \notin (\ell_3, \ell_4)$. Then the orbits of ℓ_3 or ℓ_4 under Γ are infinite. Observe that, if there is $\gamma \in \Gamma$ such that $\gamma(p_0) \in \ell_1$, the assertion follows easily. So let us assume that $p_0 \notin \Gamma \ell_1$. Since $LiG(\Omega) = k_0 < \infty$ and $\Gamma \ell_1$ is infinite, it falls out that there is $p_1 \in \mathbb{P}_{\mathbb{C}}^2 \setminus \{p_0\}$, $\ell_1, \ell_2 \in \mathcal{L}$ and a infinite set $\mathcal{C}_3 \in Gr(\mathbb{P}_{\mathbb{C}}^2)$ an infinite set such that $\{p_1\} = \bigcap \mathcal{C}_3 = \tilde{\ell}_1 \cap \tilde{\ell}_2$ and $\bigcup \mathcal{C}_3 \subset \mathbb{P}_{\mathbb{C}}^2 \setminus \Omega$. Which concludes the proof.

Let us prove (iv). On the contrary, let us assume that the vertex of $\mathcal{L}$ does not lie in a complex line, thus there is a set $C \subset \mathbb{P}_{\mathbb{C}}^2$ of vertexes of $\mathcal{L}$ such that $Card(C) = 3$. Let $\Gamma_0 = \bigcup_{x \in C} Isot(x, \Gamma)$, thus Γ_0 is a subgroup of Γ with finite index. Thus after conjugating by a projective transformation, if it is necessary, we can assume that $C = \{[e_1], [e_2], [e_3]\}$. In consequence, each element $\gamma \in \Gamma$ has a lift $\tilde{\gamma} \in SL(3, \mathbb{C})$ given by:

$$\tilde{\gamma} = \begin{pmatrix} \gamma_{11} & 0 & 0 \\ 0 & \gamma_{22} & 0 \\ 0 & 0 & \gamma_{33} \end{pmatrix}.$$

where $\gamma_{11}\gamma_{22}\gamma_{33} = 1$. Which is a contradiction.

Let us prove (v). If this is not the case, then there is a set C_1 of vertexes of $\mathcal{L}$ such that $Card(C_1) = 3$. Set $\Gamma_0 = \bigcap_{v \in C_1} Isot(v, \Gamma) \subset \Gamma$ is a subgroup of Γ with finite index fixing at least 3 points in the line ℓ which contains C_1 . Thus Γ_0 fixes each point in ℓ . By corollary (2.14), it follows

that $Eq(\Gamma) = \mathbb{P}_{\mathbb{C}}^2 \setminus \ell = \mathbb{P}_{\mathbb{C}}^2 \setminus L_0(\Gamma_0)$, which is a contradiction.

Let us prove (vi). On the contrary, let us assume that there is $\ell_0 \in Gr(\mathbb{P}_{\mathbb{C}}^2)$ an element with infinite orbit which is disjoint from the vertexes of $\mathcal{L}$. Thus $\Gamma_0 = Isot(\ell_0, \Gamma) \cap (\bigcap_{v \in V} Isot(v, \Gamma))$, is a subgroup of Γ with finite index fixing ℓ_0 and each vertex $\mathcal{L}$, here V is the set of vertex of $\mathcal{L}$. Let ℓ_V be the line which contains the vertexes of $\mathcal{L}$. Then Γ_0 leaves invariant $\ell_0 \cap \ell$. In consequence leaves invariant 3 points in ℓ . Therefore, Γ_0 leaves invariant each point in ℓ . By corollary (2.14), it follows that there is a point $q \in \mathbb{P}_{\mathbb{C}}^2$ such that $Eq(\Gamma) = \mathbb{P}_{\mathbb{C}}^2 \setminus (\ell \cup \{q\}) = \mathbb{P}_{\mathbb{C}}^2 \setminus L_0(\Gamma)$. Which is a contradiction. $\square$

Theorem 2.17. *Let $\Gamma \subset PSL(3, \mathbb{C})$ be a discrete subgroup and $\Omega \subset \mathbb{P}_{\mathbb{C}}^2$ be either $Eq(\Gamma)$, $\Omega(\Gamma)$, $\mathbb{P}_{\mathbb{C}}^2 \setminus C(\Gamma)$ or a maximal region where Γ acts properly discontinuously, then $LiG(\Omega) = 1, 2, 3, 4$ or ∞ .*

Proof. On the contrary let us assume that $4 < LiG(\Omega) < \infty$, then there is $\mathcal{L} \in LG(\Omega)$ such that $Card(\mathcal{L}) = LiG(\Omega)$. Let p_0, p_1 be the vertex of $\mathcal{L}$. Then there is $\ell \in \mathcal{L}$, disjoint from $\{p_0, p_1\}$. In consequence, previous Lemma yields that $\Gamma\ell$ is an infinite set in $Gr(\mathbb{P}_{\mathbb{C}}^2)$. Thus there is $p \in \{p_0, p_1\}$ and $(\gamma_m) \subset \Gamma$ such that $\gamma_m\ell$ is an infinite set in $Gr(\mathbb{P}_{\mathbb{C}}^2)$ and $\bigcap_{m \in \mathbb{N}} \gamma_m\ell = \{p\}$. Then $\{\gamma_m^{-1}(p) : m \in \mathbb{N}\} \subset \ell \cap \{p_0, p_1\}$. Which is a contradiction. $\square$

3. THE LIMIT SET

Lemma 3.1. *Let $\Gamma \subset PSL(3, \mathbb{C})$ be a discrete group which satisfies $LiG(\mathbb{P}_{\mathbb{C}}^2 \setminus C(\Gamma)) = 1$. If $\ell \in Gr(\mathbb{P}_{\mathbb{C}}^2)$ is the unique line in $\mathbb{P}_{\mathbb{C}}^2 \setminus C(\Gamma)$ and $\tilde{\Gamma}$ denotes the action of Γ restricted to ℓ , then $\tilde{\Gamma}$ does not contains loxodromic elements.*

Proof. On the contrary let us assume that there is $\gamma_0 \in \Gamma$ such that $\gamma_0 \mid_{\ell}$ is loxodromic. Thus Normal Jordan Form Theorem yields that there is $\tilde{\gamma}_0 \in GL(3, \mathbb{C})$ a lift of γ_0 and $\gamma \in GL(3, \mathbb{C})$ that $[[\gamma]](\ell) = \overleftrightarrow{[e_1], [e_2]}$ and $\gamma\tilde{\gamma}_0\gamma^{-1}$ is either:

$$A = \begin{pmatrix} \lambda_1 & 0 & 0 \\ 0 & \lambda_2 & 0 \\ 0 & 0 & 1 \end{pmatrix}; C = \begin{pmatrix} \lambda & 0 & 0 \\ 0 & 1 & 1 \\ 0 & 0 & 1 \end{pmatrix},$$

where $|\lambda_1| < |\lambda_2|$ and $|\lambda| \neq 1$. A simple calculation shows:

$$\mathbb{P}_{\mathbb{C}}^2 \setminus Eq(< [A] >) = \begin{cases} \overleftrightarrow{[e_2], [e_3]} \cup \{[e_1]\} & \text{if } |\lambda_2| = 1; \\ \overleftrightarrow{[e_1], [e_3]} \cup \{[e_2]\} & \text{if } |\lambda_1| = 1; \\ \overleftrightarrow{[e_2], [e_1]} \cup \overleftrightarrow{[e_2], [e_3]} & \text{if } |\lambda_2| < 1; \\ \overleftrightarrow{[e_1], [e_3]} \cup \overleftrightarrow{[e_2], [e_3]} & \text{if } |\lambda_1| < 1 < |\lambda_2|; \\ \overleftrightarrow{[e_1], [e_3]} \cup \overleftrightarrow{[e_2], [e_1]} & \text{if } |\lambda_1| > 1; \end{cases}$$

$$\mathbb{P}_{\mathbb{C}}^2 \setminus Eq(< [B] >) = \overleftrightarrow{[e_2], [e_1]} \cup \overleftrightarrow{[e_2], [e_3]}.$$

Which is not possible, since $[[\gamma]](\ell) = \overleftrightarrow{[e_1], [e_2]}$. $\square$

Lemma 3.2 (See [4]). *Let $\Gamma \subset PSL(2, \mathbb{C})$ be a group without loxodromic elements, then Γ is conjugated to a subgroup of either $SO(3)$ or $EP(\mathbb{C})$, where*

$$EP(\mathbb{C}) = \left\{ \begin{pmatrix} a & b \\ 0 & 1 \end{pmatrix} : |a| = 1 \text{ and } b \in \mathbb{C} \right\}.$$

Lemma 3.3. *Let $\Gamma \subset PSL(3, \mathbb{C})$ be a discrete group such that $LiG(\mathbb{P}_{\mathbb{C}}^2 \setminus C(\Gamma)) = 1$. If ℓ is the unique line in $C(\Gamma)$, it holds:*

- (i) *If $\Gamma|_{\ell}$ is conjugated to a group of $SO(3)$, then $LiG(Eq(\Gamma)) = 1$;*
- (ii) *If $\Gamma|_{\ell}$ is conjugated to a group of $EP(\mathbb{C})$, then either $LiG(Eq(\Gamma)) \leq 2$ or $Eq(\Gamma) = \emptyset$.*

Proof. Let us prove (i). After conjugating with a projective transformations if it is necessary, we may assume that $\ell = \overleftarrow{[e_2], [e_3]}$ and $\Gamma|_{\ell}$ is a subgroup of $SO(3)$. Now, let $(\gamma_m) \subset \Gamma$ be a sequence of distinct elements, then for each $m \in \mathbb{N}$ there is $a_m \in \mathbb{C}$, $b_m = (b_{j1}^{(m)}) \in M(2 \times 1, \mathbb{C})$, $\tilde{\gamma}_m \in GL(3, \mathbb{C})$ and $H_m = (h_{ij}^{(m)}) \in SO(3)$ such that $\tilde{\gamma}_0$ is a lift of γ_0 and

$$\tilde{\gamma}_m = \begin{pmatrix} a_m & 0 \\ b_m & H_m \end{pmatrix}.$$

On the other hand, since $SO(3)$ and $\mathbb{P}_{\mathbb{C}}^1$ are compact, we may assume that there is $a \in \mathbb{P}_{\mathbb{C}}^1$ and $H \in SO(3)$ such that $a_m \xrightarrow{m \rightarrow \infty} a$ and $H_m \xrightarrow{m \rightarrow \infty} H$.

Define $|\gamma_m| = \max\{|a_m|, |h_{ij}^{(m)}|, |b_{j1}^{(m)}|, j, i \in \{1, 2\}\}$, we must consider the following cases:

Case 1. $(|\gamma_m|)$ is unbounded. Thus there is a subsequence of $(|\gamma_m|)$, which is still denoted $(|\gamma_m|)$, such that $|\gamma_m| \xrightarrow{m \rightarrow \infty} \infty$. In consequence $(|\gamma_m|^{-1} \tilde{\gamma}_m)$ is a bounded sequence in $M(3, \mathbb{C})$, thus there is $\gamma \in M(3, \mathbb{C}) \setminus \{0\}$ and a subsequence of $(|\gamma_m|^{-1} \tilde{\gamma}_m)$, still denoted $(|\gamma_m|^{-1} \tilde{\gamma}_m)$, which satisfies $|\gamma_m|^{-1} \tilde{\gamma}_m \xrightarrow{m \rightarrow \infty} \gamma$. Since $|\gamma_m|^{-1} H_m \xrightarrow{m \rightarrow \infty} 0$, we conclude that there is $a \in \mathbb{C}$ and $b \in M(2 \times 1, \mathbb{C})$ such that:

$$\gamma = \begin{pmatrix} a & 0 \\ b & 0 \end{pmatrix}.$$

In consequence $Ker(\gamma) = \overleftarrow{[e_2], [e_3]}$.

Case 2. $(|\gamma_m|)$ is bounded. Since Γ is discrete, we conclude that $a_m \xrightarrow{m \rightarrow \infty} 0$. Moreover, $(\tilde{\gamma}_m)$ is a bounded sequence in $M(3, \mathbb{C})$, thus there is $\gamma \in M(3, \mathbb{C}) \setminus \{0\}$ and a subsequence of $(\tilde{\gamma}_m)$, still denoted $(\tilde{\gamma}_m)$, which satisfies $\tilde{\gamma}_m \xrightarrow{m \rightarrow \infty} \gamma$. Thus there are $b \in M(2 \times 1, \mathbb{C})$ such that:

$$\gamma = \begin{pmatrix} 0 & 0 \\ b & H \end{pmatrix}.$$

A straightforward calculation shows that $w_m = (1, -(H_m - a_m)^{-1}(b_m))$ is proper vector with proper value a_m . Moreover, it holds

$$w_m \xrightarrow{m \rightarrow \infty} w = (1, -H^{-1}(b))$$

and $\gamma(w) = 0$. That is $Ker([\gamma]) \subset L_0(\Gamma) \subset C(\Gamma)$.

From the previous cases, it follows that $LiG(\Gamma) = 1$.

Let us prove (ii). After conjugating with a projective transformations if it is necessary, we may assume that $\ell = \overrightarrow{[e_2], [e_3]}$ and $\Gamma|_\ell$ is a subgroup of $EP(\mathbb{C})$. Now, let $(\gamma_m) \subset \Gamma$ be a sequence of distinct elements, then for each $m \in \mathbb{N}$ there are $a_m, b_m, c_m, d_m \in \mathbb{C}$, $\theta_m \in \mathbb{R}$ and $\tilde{\gamma}_m \in GL(3, \mathbb{C})$ such that $\tilde{\gamma}_0$ is a lift of γ_0 and

$$\tilde{\gamma}_m = \begin{pmatrix} a_m & 0 & 0 \\ b_m & e^{\pi i \theta_m} & d_m \\ c_m & 0 & 1 \end{pmatrix}.$$

Define $|\gamma_m| = \max\{|a_m|, |b_m|, |c_m|, |d_m|\}$, we must consider the following cases:

Case 1. $(|\gamma_m|)$ is unbounded. Thus there is a subsequence of $(|\gamma_m|)$, which is still denoted $(|\gamma_m|)$, such that $|\gamma_m| \xrightarrow{m \rightarrow \infty} \infty$. In consequence $(|\gamma_m|^{-1} \tilde{\gamma}_m)$ is a bounded sequence in $M(3, \mathbb{C})$, thus there is $\gamma \in M(3, \mathbb{C}) \setminus \{0\}$ and a subsequence of $(|\gamma_m|^{-1} \tilde{\gamma}_m)$, still denoted $(|\gamma_m|^{-1} \tilde{\gamma}_m)$, which satisfies $|\gamma_m|^{-1} \tilde{\gamma}_m \xrightarrow{m \rightarrow \infty} \gamma$. Thus there are $a, b, c, d \in \mathbb{C}$ such that:

$$\gamma = \begin{pmatrix} a & 0 & 0 \\ b & 0 & d \\ c & 0 & 0 \end{pmatrix}.$$

A straightforward calculation shows:

$$Ker(\gamma) = \begin{cases} \overrightarrow{[e_2], [e_1]} & \text{if } d = 0; \\ \text{a line passing trough } [e_2] & \text{if } d \neq 0, a=b=0; \\ \{e_1\} & \text{in any other case} \end{cases}$$

Case 2. $(|\gamma_m|)$ is bounded. Since Γ is discrete, we conclude that $a_m \xrightarrow{m \rightarrow \infty} 0$. Moreover, $(\tilde{\gamma}_m)$ is a bounded sequence in $M(3, \mathbb{C})$, thus there is $\gamma \in M(3, \mathbb{C}) \setminus \{0\}$ and a subsequence of $(\tilde{\gamma}_m)$, still denoted $(\tilde{\gamma}_m)$, which satisfies $\tilde{\gamma}_m \xrightarrow{m \rightarrow \infty} \gamma$. Thus there are $b, c, d \in \mathbb{C}$ and $\theta \in \mathbb{R}$ such that:

$$\gamma = \begin{pmatrix} 0 & 0 & 0 \\ b & e^{\pi i \theta} & d \\ c & 0 & 1 \end{pmatrix}.$$

Set $K_m = \begin{pmatrix} e^{\pi i \theta_m} & d_m \\ 0 & 1 \end{pmatrix}$ and $K = \begin{pmatrix} e^{\pi i \theta} & d \\ 0 & 1 \end{pmatrix}$. A straightforward calculation shows that $w_m = (1, -(K_m - a_m)^{-1}(b_m, c_m))$ is proper vector with proper value a_m . Moreover, it holds

$$w_m \xrightarrow{m \rightarrow \infty} w = (1, -K^{-1}(b, c))$$

and $\gamma(w) = 0$. That is $Ker([\gamma]) \subset L_0(\Gamma) \subset C(\Gamma)$.

The previous cases, yield that either $Eq(\Gamma) = \emptyset$ or $LiG(\Gamma) \leq 2$. $\square$

Lemma 3.4. *Let $\Gamma \subset PSL(3, \mathbb{C})$ be a discrete group, such that $LiG(\mathbb{P}_{\mathbb{C}}^2 \setminus C(\Gamma)) = 2$, then:*

- (i) *There is $\tau \in PSL(3, \mathbb{C})$ a projective transformation such that each $\gamma \in \tau\Gamma\tau^{-1}$ has a lift $\tilde{\gamma} \in SL(3, \mathbb{C})$ such that $\tilde{\gamma}$ is given by:*

$$\gamma = \begin{pmatrix} \gamma_{11} & \gamma_{12} & \gamma_{13} \\ 0 & \gamma_{22} & \gamma_{23} \\ 0 & \gamma_{32} & \gamma_{33} \end{pmatrix};$$

- (ii) *Let $\gamma \in \tau\Gamma\tau^{-1}$ and $\tilde{\gamma} \in SL(3, \mathbb{C})$ a lift of γ , then there is a proper values λ of $\begin{pmatrix} \gamma_{22} & \gamma_{23} \\ \gamma_{32} & \gamma_{33} \end{pmatrix}$ such that*

$$|\gamma_{11}| \leq |\lambda|;$$

- (iii) *It holds either $LiG(Eq(\Gamma))$ or $Eq(\Gamma) = \emptyset$.*

Proof. Let us prove (i). It follows easily from lemma 2.4.

Let us prove (ii). Let $\gamma \in \tilde{\Gamma}$ and $\tilde{\gamma} \in SL(3, \mathbb{C})$ be a lift of γ , then there is $H \in GL(2, \mathbb{C})$, $a \in \mathbb{C}^*$ and $b \in M(1 \times 2, \mathbb{C})$ such that

$$\tilde{\gamma} = \begin{pmatrix} a & b \\ 0 & H \end{pmatrix}$$

Let us assume that $|a| > \max\{|\lambda| : \lambda \text{ is a proper value of } H\}$. Since every proper value of H is a proper value of $\tilde{\gamma}$, it follows that $\tilde{\gamma}$ has at least two different proper values. Thus normal Jordan form theorem yields that $\tilde{\gamma}$ is either conjugated to

$$A = \begin{pmatrix} a & 0 & 0 \\ 0 & \lambda_1 & 0 \\ 0 & 0 & \lambda_2 \end{pmatrix}; B = \begin{pmatrix} a & 0 & 0 \\ 0 & \lambda_1 & 1 \\ 0 & 0 & \lambda_1 \end{pmatrix}$$

clearly, in the respective cases, λ_1, λ_2 are proper values of H . A simple calculation shows:

$$\Lambda(<[A]>) = \begin{cases} \overline{[e_1], [e_3]} \cup \overline{[e_3], [e_2]} & |\lambda_1| < |\lambda_2|; \\ \overline{[e_1], [e_2]} \cup \overline{[e_3], [e_2]} & |\lambda_2| < |\lambda_1|; \\ \{[e_1]\} \cup \overline{[e_3], [e_2]} & |\lambda_1| = |\lambda_2| \end{cases}$$

and

$$\Lambda(< [B] >) = \overleftarrow{[e_1], [e_2]} \cup \overleftarrow{[e_3], [e_2]}$$

in any case, it follows that $LiG(\Gamma) \geq 3$. Which is a contradiction.

Let us prove (iii). Let $(\gamma_m) \subset \Gamma$ be a sequence of distinct elements and $\gamma \in SP(3, \mathbb{C})$ such that $\gamma_m \xrightarrow{m \rightarrow \infty} \gamma$. Thus for each $m \in n\mathbb{N}$ there is $\tilde{\gamma}_m \in SL(3, \mathbb{C})$, $\gamma_{11}^{(m)} \in \mathbb{C}^*$, $\gamma_{12}^{(m)} \in M(1 \times 2, \mathbb{C})$ and $H_m \in GL(2, \mathbb{C})$ such that $\tilde{\gamma}_m$ is a lift of γ_m and

$$\tilde{\gamma}_m = \begin{pmatrix} \gamma_{11}^{(m)} & \gamma_{12}^{(m)} \\ 0 & H_m \end{pmatrix}.$$

Set $\|H_m\| = \max\{H_m(x) : |x| = 1\}$. Thus $(\|H_m\|^{-1}H_m)$ is sequence of unitary operators, thus we can assume that there is $H \in GL(2, \mathbb{C})$ a unitary operator such that $\|H_m\|^{-1}H_m \xrightarrow{m \rightarrow \infty} H$, point wise. On the other hand, part (ii) of the present lemma yields $|\gamma_{11}^{(m)}| \leq \|H_m\|$. Then we can assume that there is $\gamma_{11} \in \mathbb{C}$ such that $\gamma_{11}^{(m)}\|H_m\|^{-1} \xrightarrow{m \rightarrow \infty} \gamma_{11}$. Set $\gamma_{12}^{(m)} = (a_m, b_m)$ and $|\gamma_{12}(m)| = \max\{|a_m|, |b_m|\}$. And consider the following cases:

Case 1.- The sequence $(\|H_m\|^{-1}|\gamma_{12}^{(m)}|)$ is unbounded. In this case, we may assume that $\|H_m\|^{-1}|\gamma_{12}^{(m)}| \xrightarrow{m \rightarrow \infty} \infty$. Thus a simple calculation shows:

$$|\gamma_{12}^{(m)}|^{-1}\|H_m\|\tilde{\gamma}_m \xrightarrow{m \rightarrow \infty} \begin{pmatrix} 0 & \gamma_{12} \\ 0 & 0 \end{pmatrix}.$$

which shows that $Ker(\gamma)$ is a complex line passing trough $[e_1]$.

Case 2.- The sequence $(\|H_m\|^{-1}|\gamma_{12}^{(m)}|)$ is bounded. In this case, we may assume that there is $\gamma_{12} \in M(1 \times 2, \mathbb{C})$ such that $\|H_m\|^{-1}\gamma_{12}^{(m)} \xrightarrow{m \rightarrow \infty} \gamma_{12}$. A simple calculation shows:

$$\|H_m\|^{-1}\tilde{\gamma}_m \xrightarrow{m \rightarrow \infty} \begin{pmatrix} \gamma_{11} & \gamma_{12} \\ 0 & H \end{pmatrix}.$$

Now is straightfoward to check that $Ker(\gamma) = [e_1]$ is either $[e_1]$, a line passing trough $[e_1]$ or a point in $\mathbb{P}_{\mathbb{C}}^2$. Let us assume that $Ker(\gamma)$ is a point $*$. In thus case there is $\tilde{\ell} \in Gr(\mathbb{P}_{\mathbb{C}}^2)$ such that $* \in \ell \subset C(\Gamma)$. It can be shown that, see [?], there is a complex line $\hat{\ell} \in Gr(\mathbb{P}_{\mathbb{C}}^2)$ and subsequence of (γ_m) , still denoted (γ_m) , such that

$$(3.1) \quad \gamma_m^{-1} \xrightarrow{m \rightarrow \infty} * \text{ uniformly on compact sets of } \mathbb{P}_{\mathbb{C}}^2 \setminus \hat{\ell}.$$

Let $\check{\ell} \in Gr(\mathbb{P}_{\mathbb{C}}^2)$ be such that $\check{\ell} \subset C(\Gamma)$ and $\check{\ell} \neq \hat{\ell}$. Thus equation 3.1, yields $\gamma_m^{-1}(\check{\ell}) \xrightarrow{m \rightarrow \infty} \overleftarrow{*, [e_1]}$. Which clearly concludes the proof. $\square$

From the previous lemmas is straightfoward to check

Corollary 3.5. *Let $\Gamma \subset PSL(3, \mathbb{C})$ be a complex Kleinian group such that $LiG(\Omega(\Gamma)) \geq 3$, then $LiG(\mathbb{P}_{\mathbb{C}}^2 \setminus C(\Gamma)) > 3$.*

Now Theorem 0.3 follows easily.

4. EXAMPLES

4.1. A group With infinite Lines in General Position. Let $\Gamma \subset PU(1, 2)$ be a discrete group which acts with compact quotient on $\mathbb{H}_{\mathbb{C}}^2$, see [3], then it can be proved that, see [9], $\Omega(\Gamma) = \mathbb{H}_{\mathbb{C}}^2$ and $LiG(\Omega) = \infty$.

4.2. A group With 4 Lines in General Position. Let M be the matrix $M = \begin{pmatrix} 3 & 5 \\ -5 & 8 \end{pmatrix}$. A straightforward calculation shows, that M has eigenvalues

$$\alpha_{\pm} = \frac{-5 \pm \sqrt{21}}{2},$$

with respective eigenvectors:

$$v_+ = \left(1, \frac{-11 + \sqrt{21}}{10} \right); v_- = \left(\frac{-11 + \sqrt{21}}{10}, 1 \right).$$

Let $\Gamma_M^{\times} \subset PSL(3, \mathbb{C})$ be the group generated by:

$$\begin{aligned} \gamma_0(w, z) &= (\alpha_+ w, \alpha_- z); \\ \gamma_1(w, z) &= (w + 1, z + \frac{-11 + \sqrt{21}}{10}); \\ \gamma_2(w, z) &= (w + \frac{-11 + \sqrt{21}}{10}, z + 1); \\ \gamma_3(w, z) &= (z, w). \end{aligned}$$

Then it is easily seen that $\Omega(\Gamma_A^{\times}) = \bigcup_{j,j=0,1} (\mathbb{H}^{(-1)^i} \times \mathbb{H}^{(-1)^j})$ and $LiG(\Omega(\Gamma_A^{\times})) = 4$. Moreover, one can check that this is "essentially" the only way to construct examples of groups $\Gamma \in PSL(3, \mathbb{C})$ with $LiG(\Omega(\Gamma)) = 4$, see [2].

4.3. A group With Three Lines in General Position and infinite Lines. Let $\Gamma \subset PSL(2, \mathbb{C})$ be a non-elementary kleinian group and $G \subset \mathbb{C}^*$ be a discrete subgroup and for each $g \in \mathbb{C}^*$ set $i_g : SL(2, \mathbb{C}) \rightarrow SL(3, \mathbb{C})$ given by:

$$i(h) = \begin{pmatrix} gh & 0 \\ 0 & g^{-2} \end{pmatrix}.$$

Then $S(\Gamma, G)$ is to be defined as:

$$S(\Gamma, G) = \{i_g(\gamma) : g \in G \text{ \& } [[\gamma]] \in \Gamma\}.$$

Then, one can check $\Omega(S(\Gamma, G)) = \Omega(\Gamma) \times \mathbb{C}^*$, $LiG(\Omega(S(\Gamma, G))) = 3$ and $Li(\Omega(S(\Gamma, G))) = \infty$

4.4. A group With Three Lines in General Position. Let $a \in \mathbb{C}^*$ and $M_a, B \in SL(3, \mathbb{C})$ given by:

$$M_a = \begin{pmatrix} a & 0 & 0 \\ 0 & a & 0 \\ 0 & 0 & a^{-2} \end{pmatrix}; B = \begin{pmatrix} 0 & 0 & 1 \\ 1 & 0 & 0 \\ 0 & 1 & 0 \end{pmatrix}$$

Let $\Gamma_a \subset PSL(3, \mathbb{C})$ be the group generated by M_a and B , thus one can check, see [4], $\Omega(\Gamma_a) = \mathbb{C}^* \times \mathbb{C}^*$ and $LiG(\Omega(\Gamma_a)) = Li(\Omega(\Gamma_a)) = 3$.

4.5. A group With Two Lines in General Position And Infinite Lines. Let $M \in SL(3, \mathbb{Z})$, also let $\alpha, \beta, \bar{\beta}$ with $\alpha > 1$, $\beta \neq \bar{\beta}$ be its eigenvalues. Choose a real eigenvector (a_1, a_2, a_3) belonging to α and an eigenvector (b_1, b_2, b_3) belonging to β . Now let G_M be the group generated by:

$$\begin{aligned} \gamma_0(w, z) &= (\alpha w, \beta z), \\ \gamma_i(w, z) &= (w + a_i, z + b_i); i = 1, 2, 3. \end{aligned}$$

Then it is easily seen that $\Omega(G_M) = \bigcup_{j=0,1} (\mathbb{H}^{(-1)^j} \times \mathbb{C})$, $Li(\Omega(G_M)) = \infty$, $LiG(\Omega(G_M)) = 2$ and $Eq(G_M) = \emptyset$.

4.6. A group With Two Lines in General Position. Let $\Gamma \subset PSL(3, \mathbb{C})$ be the group induce by the matrix:

$$\gamma = \begin{pmatrix} a & 0 & 0 \\ 1 & b & 0 \\ 0 & 0 & c \end{pmatrix},$$

where $|a| < |b| < |c|$ and $abc = 1$. Thus it is easily seen, see [10], that $\Omega(\Gamma) = \mathbb{C} \times \mathbb{C}^*$ and in consequence $LiG(\Omega(\Gamma)) = Li(\Omega(\Gamma)) = 2$.

4.7. A group With One Line in General Position. Let $v_1, v_2, v_3, v_4 \in \mathbb{C}^2$ be points which are $\mathbb{R}$ -linearly independent. Let g_i , $i \in \{1, 2, 3, 4\}$, be the translation induced by v_i , then $\Gamma = \langle g_1, \dots, g_4 \rangle$ is a group isomorphic to $\mathbb{Z}^4$ which satisfy $\Omega(\Gamma) = \mathbb{C}^2$ and $LiG(\Omega(\Gamma)) = 1$.

The authors are grateful to Professor José Seade for his stimulating conversations and encouragement. Part of this research was done while the authors were visiting the IMATE-UNAM campus Cuernavaca and the FMAT of the UADY. Also, during this time, the second author was in a postdoctoral year at IMPA, and they are grateful to these institutions and its people, for their support and hospitality.

REFERENCES

1. W. Barrera, A. Cano, J. P. Navarrete, *The limit set of discrete subgroups of $PSL(3, \mathbb{C})$* , preprint 2009, <http://arxiv.org/abs/1002.0021>.
2. W. Barrera, A. Cano, J. P. Navarrete, *Two dimensional Complex Kleinian Groups With Four Complex Lines in General Position in its Limit Set*, preprint 2010, <http://arxiv.org/abs/1001.5222>.
3. P. Deligne, G. D. Mostow. *Commensurabilities among lattices in $PU(1, n)$* . Ann. Math. Study **132** (1993), Princeton Univ. Press.

4. A. Cano, *On Discrete Subgroups of automorphism of $\mathbb{P}_{\mathbb{C}}^2$* , preprint 2009, <http://arxiv.org/abs/0806.1336>.
5. A. Cano, J.-P. Navarrete, J. Seade. *Complex Kleinian Groups*. Monograph to be published, <http://www.matcuer.unam.mx/~jseade/Libro-Klein.pdf>.
6. A. Cano, J. Seade. *On the equicontinuity region of discrete subgroups of $PU(n, 1)$* , JGA, Vol 20, No. 2, 2010.
7. S. Kobayashi, *Hyperbolic Manifolds and Holomorphic Mappings*, 148 pp. Marcel Dekker, 1970.
8. R. S. Kulkarni. *Groups with Domains of Discontinuity*, Math. Ann. **237**, 253-272 (1978).
9. J.-P. Navarrete. *On the limit set of discrete subgroups of $PU(2, 1)$* . Geom. Dedicata 122, 1-13 (2006).
10. J.-P. Navarrete. *The Trace Function and Complex Kleinian Groups in $\mathbb{P}_{\mathbb{C}}^2$* . Int. Jour. of Math. Vol. 19, No. 7 (2008).

WALDEMAR BARRERA: UNIVERSIDAD AUTÓNOMA DE YUCATÁN FACULTAD DE MATEMÁTICAS, ANILLO PERIFÉRICO NORTE TABLAJE CAT 13615 CHUBURNÁ HIDALGO, MÉRIDA YUCATÁN. MÉXICO., ANGEL CANO: INSTITUTO DE MATEMÁTICA PURA E APLICADA (IMPA), RIO DE JANEIRO, BRAZIL., JUAN PABLO NAVARRETE: UNIVERSIDAD AUTÓNOMA DE YUCATÁN FACULTAD DE MATEMÁTICAS, ANILLO PERIFÉRICO NORTE TABLAJE CAT 13615 CHUBURNÁ HIDALGO, MÉRIDA YUCATÁN. MÉXICO.,

E-mail address: bvargas@uady.mx, angel@impa.br, jp.navarrete@uady.mx